GAD-7 Anxiety

Over the <u>last two weeks</u> , how often have you been bothered by the following problems?	Not at all	Several days	More than half the days	Nearly every day
Feeling nervous, anxious, or on edge	0	1	2	3
Not being able to stop or control worrying	0	1	2	3
Worrying too much about different things	0	1	2	3
4. Trouble relaxing	0	1	2	3
5. Being so restless that it is hard to sit still	0	1	2	3
Becoming easily annoyed or irritable	0	1	2	3
 Feeling afraid, as if something awful might happen 	0	1	2	3
Column totals	+		+	· =
Total score				
If you checked any problems, how difficult have they made it for you to do your work, take care of things at home, or get along with other people?				

Source: Primary Care Evaluation of Mental Disorders Patient Health Questionnaire (PRIME-MD-PHQ). The PHQ was
developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke, and colleagues. For research information, contact Dr.
Spitzer at ris8@columbia.edu. PRIME-MD® is a trademark of Pfizer Inc. Copyright© 1999 Pfizer Inc. All rights reserved.
Reproduced with permission

Somewhat difficult

Very difficult

Extremely difficult

Scoring GAD-7 Anxiety Severity

This is calculated by assigning scores of 0, 1, 2, and 3 to the response categories, respectively, of "not at all," "several days," "more than half the days," and "nearly every day." GAD-7 total score for the seven items ranges from 0 to 21.

0-4: minimal anxiety

5-9: mild anxiety

Not difficult at all

10–14: moderate anxiety

15-21: severe anxiety

Mood Disorder Questionnaire (MDQ)

Name: Date:		
Instructions: Check () the answer that best applies to you. Please answer each question as best you can.	Yes	No
1. Has there ever been a period of time when you were not your usual self and		
you felt so good or so hyper that other people thought you were not your normal self or you were so hyper that you got into trouble?		
you were so irritable that you shouted at people or started fights or arguments?		
you felt much more self-confident than usual?		
you got much less sleep than usual and found you didn't really miss it?		
you were much more talkative or spoke faster than usual?		
thoughts raced through your head or you couldn't slow your mind down?		
you were so easily distracted by things around you that you had trouble concentrating or staying on track?		
you had much more energy than usual?		
you were much more active or did many more things than usual?		
you were much more social or outgoing than usual, for example, you telephoned friends in the middle of the night?		
you were much more interested in sex than usual?		
you did things that were unusual for you or that other people might have thought were excessive, foolish, or risky?		
spending money got you or your family in trouble?		
2. If you checked YES to more than one of the above, have several of these ever happened during the same period of time? Please check 1 response only.	\circ	
3. How much of a problem did any of these cause you — like being able to work; having family, money, or legal troubles; getting into arguments or fights? Please check 1 response only.		
No problem Minor problem Moderate problem Serious problem		
4. Have any of your blood relatives (ie, children, siblings, parents, grandparents, aunts, uncles) had manic-depressive illness or bipolar disorder?		
5. Has a health professional ever told you that you have manic-depressive illness or bipolar disorder?		

This questionnaire should be used as a starting point. It is not a substitute for a full medical evaluation. Bipolar disorder is a complex illness, and **an accurate, thorough diagnosis can only be made through a personal evaluation by your doctor.**

This instrument is designed for screening purposes only and is not to be used as a diagnostic tool.

How to Use

The questionnaire takes less than 5 minutes to complete. Patients simply check the yes or no boxes in response to the questions. The last question pertains to the patient's level of functional impairment. The physician, nurse, or medical staff assistant then scores the completed questionnaire.

How to Score

Further medical assessment for bipolar disorder is clearly warranted if patient:

 Answers Yes to 7 or more of the events in question #1

AND

• Answers *Yes* to question #2

AND

• Answers Moderate problem or Serious problem to question #3

PATIENT HEALTH QUESTIONNAIRE-9 (PHQ-9)

by any of the following pro (Use "✔" to indicate your an		Not at all	Several days	More than half the days	Nearly every day
1. Little interest or pleasure i	n doing things	0	1	2	3
2. Feeling down, depressed,	or hopeless	0	1	2	3
3. Trouble falling or staying a	asleep, or sleeping too much	0	1	2	3
4. Feeling tired or having littl	e energy	0	1	2	3
5. Poor appetite or overeating	g	0	1	2	3
Feeling bad about yoursel have let yourself or your factors	f — or that you are a failure or amily down	0	1	2	3
7. Trouble concentrating on newspaper or watching te		0	1	2	3
noticed? Or the opposite	wly that other people could have — being so fidgety or restless g around a lot more than usual	0	1	2	3
Thoughts that you would be yourself in some way	oe better off dead or of hurting	0	1	2	3
	FOR OFFICE CO	ding <u>0</u> +			
				Total Score	:
	blems, how <u>difficult</u> have these t home, or get along with other		ade it for	you to do y	your
Not difficult at all □	Somewhat difficult □	Very difficult □		Extremely difficult □	

PCL-C

<u>INSTRUCTIONS</u>: Below is a list of problems and complaints that people sometimes have in response to stressful life experiences. Please read each one carefully, then circle one of the numbers to the right to indicate how much you have been bothered by that problem <u>in the past month</u>.

		Not at all	A little bit	Moderately	Quite a bit	Extremely
1.	Repeated, disturbing <i>memories, thoughts,</i> or <i>images</i> of a stressful experience from the past?	1	2	3	4	5
2.	Repeated, disturbing <i>dreams</i> of a stressful experience from the past?	1	2	3	4	5
3.	Suddenly <i>acting</i> or <i>feeling</i> as if a stressful experience <i>were happening again</i> (as if you were reliving it)?	1	2	3	4	5
4.	Feeling <i>very upset</i> when <i>something reminded you</i> of a stressful experience from the past?	1	2	3	4	5
5.	Having <i>physical reactions</i> (e.g., heart pounding, trouble breathing, sweating) when <i>something reminded you</i> of a stressful experience from the past?	1	2	3	4	5
6.	Avoiding <i>thinking about</i> or <i>talking about</i> a stressful experience from the past or avoiding <i>having feelings</i> related to it?	1	2	3	4	5
7. ——	Avoiding <i>activities</i> or <i>situations</i> because <i>they reminded you</i> of a stressful experience from the past?	1	2	3	4	5
8.	Trouble <i>remembering important parts</i> of a stressful experience from the past?	1	2	3	4	5
9.	Loss of interest in activities that you used to enjoy?	1	2	3	4	5
10.	Feeling distant or cut off from other people?	1	2	3	4	5
11.	Feeling <i>emotionally numb</i> or being unable to have loving feelings for those close to you?	1	2	3	4	5
12.	Feeling as if your <i>future</i> will somehow be <i>cut short</i> ?	1	2	3	4	5
13.	Trouble falling or staying asleep?	1	2	3	4	5
14.	Feeling irritable or having angry outbursts?	1	2	3	4	5
15.	Having difficulty concentrating?	1	2	3	4	5
16.	Being "super-alert" or watchful or on guard?	1	2	3	4	5
17.	Feeling <i>jumpy</i> or easily startled?	1	2	3	4	5